

**Table 1**  
**Survey Location Covertypes**  
**Lake Success Business Park**  
**Bridgeport, CT**

Survey Location	Covertypes			
	Wetland <sup>1</sup>	Forest	Successional	Open/Former Industrial
LSBP01	X	X		<b>X</b>
LSBP02	<b>X</b>	X		
LSBP03	X	<b>X</b>		
LSBP04				X
LSBP05		X		
LSBP06	<b>X</b>	X	X	
LSBP07				X
LSBP08		X		
LSBP09		X		<b>X</b>
LSBP10		X		<b>X</b>
LSBP11	<b>X</b>		X	
LSBP12	X			
LSBP13		X		
LSBP14	X	<b>X</b>		X
LSBP15		X		
LSBP16		X		
LSBP17		X		
LSBP18	X	<b>X</b>		
LSBP19				X
LSBP20		X		

**Notes:**

Most plots were located in areas of more than one covertypes.

Bolded "X" indicates primary covertypes at locations adjacent to more than one covertypes.

<sup>1</sup>Includes open water (Lake Success)

**Table 2**  
**Wildlife Species Observed or Known to Occur On-Site**  
**Lake Success Business Park**  
**Bridgeport, CT**

Common Name	Scientific Name	% of Plots Observed	Average # of Individuals/day	Average # of Individuals/plot
<b>Avian Species Observed During Surveys</b>				
American Robin	<i>Turdus migratorius</i>	100	23.8	1.9
Common Grackle	<i>Quiscalus quiscula</i>	90	37.8	3.6
Blue Jay	<i>Cyanocitta cristata</i>	90	13.8	1.7
Red-bellied Woodpecker	<i>Melanerpes carolinus</i>	85	10.8	1.2
Baltimore Oriole	<i>Icterus galbula</i>	80	9.3	1.4
Gray Catbird	<i>Dumetella carolinensis</i>	80	14.0	1.4
Eastern Towhee	<i>Pipilo erythrophthalmus</i>	75	6.8	1.1
Yellow Warbler	<i>Dendroica petechia</i>	70	13.5	1.5
Tufted Titmouse	<i>Baeolophus bicolor</i>	70	6.0	1.2
Mourning Dove	<i>Zenaida macroura</i>	65	13.5	1.9
Northern Cardinal	<i>Cardinalis cardinalis</i>	65	6.3	1.1
Eastern Wood-Pewee	<i>Contopus virens</i>	65	6.0	1.0
Red-winged Blackbird	<i>Agelaius phoeniceus</i>	60	26.3	2.7
Chimney Swift	<i>Chaetura pelagica</i>	60	5.5	1.4
Great Crested Flycatcher	<i>Myiarchus crinitus</i>	60	7.0	1.3
Common Yellowthroat	<i>Geothlypis trichas</i>	60	6.5	1.1
Northern Flicker	<i>Colaptes auratus</i>	60	3.8	1.0
Northern Mockingbird	<i>Mimus polyglottos</i>	55	6.5	1.1
Brown-headed Cowbird	<i>Molothrus ater</i>	45	7.5	1.6
White-breasted Nuthatch	<i>Sitta carolinensis</i>	45	3.8	1.2
Brown Thrasher	<i>Toxostoma rufum</i>	45	3.0	1.0
Wood Thrush	<i>Hylocichla mustelina</i>	40	3.3	1.2
American Goldfinch	<i>Carduelis tristis</i>	35	3.5	1.8
Warbling Vireo	<i>Vireo gilvus</i>	35	2.8	1.2
Black-and-white Warbler	<i>Mniotilta varia</i>	35	2.5	1.1
Song Sparrow	<i>Melospiza melodia</i>	35	5.3	1.1
Cedar Waxwing	<i>Bombycilla cedrorum</i>	30	4.5	2.6
American Crow	<i>Corvus brachyrhynchos</i>	30	4.3	2.1
Northern Rough-Winged Swallow	<i>Stelgidopteryx serripennis</i>	25	3.5	1.8
Herring Gull	<i>Larus argentatus</i>	25	2.5	1.7
Red-tailed Hawk	<i>Buteo jamaicensis</i>	25	2.0	1.6
Eastern Phoebe	<i>Sayornis phoebe</i>	25	4.3	1.4
Red-eyed Vireo	<i>Vireo olivaceus</i>	25	2.0	1.1
Blue-winged Warbler	<i>Vermivora pinus</i>	25	2.0	1.0
Black-crowned Night-Heron	<i>Nycticorax nycticorax</i>	25	1.8	1.0
European Starling	<i>Sturnus vulgaris</i>	25	1.5	1.0
Canada Goose	<i>Branta canadensis</i>	20	12.0	5.3
Eastern Kingbird	<i>Tyrannus tyrannus</i>	20	2.0	1.1
Field Sparrow	<i>Spizella pusilla</i>	20	1.8	1.0
House Wren	<i>Troglodytes aedon</i>	20	1.3	1.0
Wood Duck	<i>Aix sponsa</i>	15	11.0	6.3
House Sparrow	<i>Passer domesticus</i>	15	3.8	3.0
Unidentified Flycatcher	NA	15	1.0	1.3
Rock Dove	<i>Columba livia</i>	15	1.0	1.3
Indigo Bunting	<i>Passerina cyanea</i>	15	1.3	1.0
Black-capped Chickadee	<i>Poecile atricapilla</i>	15	1.0	1.0
Great Blue Heron	<i>Ardea herodias</i>	15	1.0	1.0
Mute Swan	<i>Cygnus olor</i>	10	2.5	2.0
Tree Swallow	<i>Tachycineta bicolor</i>	10	1.0	2.0

**Table 2**  
**Wildlife Species Observed or Known to Occur On-Site**  
**Lake Success Business Park**  
**Bridgeport, CT**

Common Name	Scientific Name	% of Plots Observed	Average # of Individuals/day	Average # of Individuals/plot
Barn Swallow	<i>Hirundo rustica</i>	10	0.8	1.0
Mallard Duck	<i>Anas platyrhynchos</i>	10	0.5	1.0
Wild Turkey	<i>Meleagris gallopavo</i>	10	0.5	1.0
Unidentified Woodpecker	<i>Picoides sp.</i>	10	0.5	1.0
Killdeer	<i>Charadrius vociferus</i>	5	0.8	3.0
Barred Owl	<i>Strix varia</i>	5	0.5	2.0
Carolina Wren	<i>Thryothorus ludovicianus</i>	5	0.5	2.0
Hairy Woodpecker	<i>Picoides villosus</i>	5	0.5	1.0
Belted Kingfisher	<i>Ceryle alcyon</i>	5	0.3	1.0
Blackburnian Warbler	<i>Dendroica fusca</i>	5	0.3	1.0
Green Heron	<i>Butorides virescens</i>	5	0.3	1.0
Orchard Oriole	<i>Icterus spurius</i>	5	0.3	1.0
Rose-breasted Grosbeak	<i>Pheucticus ludovicianus</i>	5	0.3	1.0
Scarlet Tanager	<i>Piranga olivacea</i>	5	0.3	1.0
Turkey Vulture	<i>Cathartes aura</i>	5	0.3	1.0
Veery	<i>Catharus fuscescens</i>	5	0.3	1.0
<b>Other Avian Species Known to Occur or Observed Directly</b>				
Black-throated Green Warbler	<i>Dendroica virens</i>		--	
Monk Parakeet	<i>Myiopsitta monachus</i>		--	
Osprey	<i>Pandion haliaetus</i>		--	
<b>Fish Known to Occur or Observed Directly</b>				
Bluegill sunfish	<i>Lepomis macrochirus</i>		--	
Chain pickerel	<i>Esox niger</i>		--	
Common carp	<i>Cyprinus carpio</i>		--	
Crappie	<i>Pomoxis sp.</i>		--	
Largemouth bass	<i>Micropterus salmoides</i>		--	
<b>Herptiles Known to Occur or Observed Directly</b>				
Bullfrog	<i>Rana catesbeiana</i>		--	
Spotted salamander	<i>Ambystoma maculatum</i>		--	
Spring peeper	<i>Pseudacris crucifer</i>		--	
Snapping turtle	<i>Chelydra serpentina serpentina</i>		--	
Turtle	NA		--	
<b>Mammals Known to Occur or Observed Directly</b>				
Eastern chipmunk	<i>Tamias striatus</i>		--	
Eastern cottontail rabbit	<i>Sylvilagus floridanus</i>		--	
Eastern gray squirrel	<i>Sciurus carolinensis</i>		--	
Fisher	<i>Martes pennanti</i>		--	
Raccoon	<i>Procyon lotor</i>		--	
Striped skunk	<i>Mephitis mephitis</i>		--	
White-tailed deer	<i>Odocoileus virginianus</i>		--	

**Notes:**

-- = Data not available for calculation

Data for Avian Species Observed During Survey are sorted first by the percent of plots observed followed by the average number of individuals per day.

The percent of plots observed is calculated as the number of plots the species was observed in all four surveys divided by 20 plots.

Average per day is calculated as the total number of birds observed during all four surveys divided by the number of surveys (4).

Average per plot is calculated as the total number of birds observed during all four surveys divided by the number of plots the species was observed.